

ULTRADISCRETIZATION OF THE TZITZEICA EQUATION

RYOGO HIROTA

*Emeritus Professor, Waseda University,
3-4-1, Ohkubo, Shinjuku-ku, Tokyo 169-8555, Japan
e-mail: roy@spn1.speednet.ne.jp*

and DAISUKE TAKAHASHI

*Department of Mathematical Sciences, Waseda University,
3-4-1, Ohkubo, Shinjuku-ku, Tokyo 169-8555, Japan
e-mail:daisuket@waseda.jp*

(Received)

Abstract

The trilinear form of the discrete Tzitzeica equation by Schief is found to be a discrete Toda molecule equation with a special boundary condition. Based on this fact, a higher order discrete Tzitzeica equation and a ultradiscrete Tzitzeica equation are obtained.

1991 *Mathematical Subject Classification.* 35Q51

1 Introduction

Tzitzeica [1, 2] obtained the equation

$$(\log h)_{xy} = h - h^{-2} \tag{1}$$

as the compatibility condition of the Gauss equation

$$\begin{aligned} r_{xx} &= (h_x r_x + \lambda r_y)/h, \\ r_{xy} &= hr, \\ r_{yy} &= (h_y r_y + \lambda^{-1} r_x)/h, \end{aligned}$$

in connection with an *affine sphere*.

Kaptsov and Shan'ko [3] transformed the Tzitzeica equation, using the dependent variable transformation

$$v = 1 - 2(\log \tau)_{xy},$$

into

$$(1 - 2(\log \tau)_{x,y})^2 \{(\log(\tau^2 - 2\tau\tau_{xy} + 2\tau_x\tau_y))_{xy} - 1\} + 1 = 0, \quad (2)$$

whose numerator is a trilinear equation. They have obtained N-soliton solutions to the trilinear equation.

On the other hand, Schief [4, 5] have obtained an integrable discrete version of the Tzitzeica equation as the compatibility condition of the discrete Gauss equation,

$$\begin{aligned} \mathbf{r}_{11} - \mathbf{r}_1 &= \alpha(\mathbf{r}_1 - \mathbf{r}) + \beta(\mathbf{r}_{12} - \mathbf{r}_1), \\ \mathbf{r}_{12} + \mathbf{r} &= H(\mathbf{r}_1 + \mathbf{r}_2), \\ \mathbf{r}_{22} - \mathbf{r}_2 &= \gamma(\mathbf{r}_2 - \mathbf{r}) + \delta(\mathbf{r}_{12} - \mathbf{r}_2). \end{aligned}$$

Schief has transformed the discrete Tzitzeica equation into the trilinear form

$$\begin{aligned} &\begin{vmatrix} \tau(m, n) & \tau(m, n+1) & \tau(m, n+2) \\ \tau(m+1, n) & \tau(m+1, n+1) & \tau(m+1, n+2) \\ \tau(m+2, n) & \tau(m+2, n+1) & \tau(m+2, n+2) \end{vmatrix} \\ &= ab \tau(m+1, n+1)^3, \quad a, b \text{ being parameters} \end{aligned} \quad (3)$$

which is reduced to the following trilinear equation in the continuous limit

$$\begin{vmatrix} \tau & \tau_x & \tau_{xx} \\ \tau_y & \tau_{xy} & \tau_{xxy} \\ \tau_{yy} & \tau_{xyy} & \tau_{xxyy} \end{vmatrix} = c \tau^3, \quad c \text{ being a parameter.} \quad (4)$$

We shall transform the Tzitzeica equation(1) into the bilinear form and find a relation between the trilinear equations (4) and (2). Let $h = G/F$. Then we have

$$\frac{\partial^2}{\partial x \partial y} \log h = \frac{G_{xy}G - G_x G_y}{G^2} - \frac{F_{xy}F - F_x F_y}{F^2}.$$

Here we introduce the bilinear operators D_x and D_y operating on an ordered pair of F and G

$$D_x^n F \cdot G = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^n F(x)G(x') \Big|_{x'=x}$$

$$D_x^n D_y^m F \cdot G = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^n \left(\frac{\partial}{\partial y} - \frac{\partial}{\partial y'} \right)^m F(x, y)G(x', y') \Big|_{x'=x, y'=y}$$

which give

$$D_x D_y F \cdot F = 2(F_{xy}F - F_x F_y)$$

$$D_x D_y G \cdot G = 2(G_{xy}G - G_x G_y).$$

Accordingly the Tzitzeica equation(1) is transformed into

$$\frac{D_x D_y G \cdot G - 2\alpha G^2 + 2F^2}{2G^2} - \frac{D_x D_y F \cdot F - 2\alpha F^2 + 2GF}{2F^2} = 0,$$

where α being an arbitrary parameter. Accordingly the Tzitzeica equation(1) is decoupled into the bilinear equations

$$D_x D_y G \cdot G = 2(\alpha G^2 - F^2), \quad (5)$$

$$D_x D_y F \cdot F = 2(\alpha F^2 - GF). \quad (6)$$

Let us introduce new dependent variables τ_1 and express F and G as follows

$$F = \tau_1^2, \quad (7)$$

$$G = \alpha \tau_1^2 - D_x D_y \tau_1 \cdot \tau_1. \quad (8)$$

Equation(6) is satisfied by this choice of F and G , because of the identity

$$D_x D_y \tau_1^2 \cdot \tau_1^2 = 2\tau_1^2 D_x D_y \tau_1 \cdot \tau_1.$$

Let $G = \tau_2$. Then eqs.(8) and (5) are written as

$$D_x D_y \tau_1 \cdot \tau_1 = \alpha \tau_1^2 - \tau_2, \quad (9)$$

$$D_x D_y \tau_2 \cdot \tau_2 = 2(\alpha \tau_2^2 - \tau_1^4). \quad (10)$$

Equations (9) and (10) are reduced to the trilinear equations (2) and (4) for $\alpha = 1$ and $\alpha = 0$, respectively, by eliminating τ_2 .

2 Toda molecule equation

Equations (9) and (10) for $\alpha = 0$ remind us of the Toda molecule equation expressed with the bilinear form

$$D_x D_y \tau_n \cdot \tau_n = 2\tau_{n+1}\tau_{n-1}, \quad (11)$$

for $n = 1, 2, \dots, N$. For $N = 2$ we have

$$D_x D_y \tau_1 \cdot \tau_1 = 2\tau_2\tau_0, \quad (12)$$

$$D_x D_y \tau_2 \cdot \tau_2 = 2\tau_3\tau_1, \quad (13)$$

which become eqs. (9) and (10) for $\alpha = 0$ by choosing the boundary condition

$$\tau_0 = -1/2, \quad \tau_3 = -\tau_1^3. \quad (14)$$

The bilinear equation (11) is transformed into the ordinary form [6]

$$\frac{\partial^2}{\partial x \partial y} \log V_n = V_{n+1} - 2V_n + V_{n-1}, \quad (15)$$

for $n = 1, 2, \dots, N$, through the dependent variable transformation

$$V_n = \frac{\partial^2}{\partial x \partial y} \log \tau_n = \frac{\tau_{n+1}\tau_{n-1}}{\tau_n^2}. \quad (16)$$

Accordingly eqs.(12) and (13) with the boundary condition (14) are transformed into

$$\frac{\partial^2}{\partial x \partial y} \log V_1 = V_2 - 2V_1, \quad (17)$$

$$\frac{\partial^2}{\partial x \partial y} \log V_2 = 4V_1 - 2V_2. \quad (18)$$

Adding twice of eq.(17) to eq.(18) we obtain

$$\frac{\partial^2}{\partial x \partial y} \log(V_1^2 V_2) = 0, \quad (19)$$

which is consistent with

$$V_2 = -\frac{1}{4}V_1^{-2}, \quad (20)$$

given by eqs. (14) and (16).

Substituting eq. (20) into eq.(17) we obtain

$$\frac{\partial^2}{\partial x \partial y} \log V_1 = -\frac{1}{4}V_1^{-2} - 2V_1. \quad (21)$$

Let

$$V_1 = -\frac{h}{2}. \quad (22)$$

Then eq.(21) becomes the Tzitzeica equation

$$\frac{\partial^2}{\partial x \partial y} \log h = h - h^{-2}. \quad (23)$$

Hence we have shown that the Tzitzeica equation is equivalent to the Toda molecule equation with the special boundary condition. The Toda molecule equation is discretized in [7]. Hence the discrete Tzitzeica equation could be obtained by using the discrete Toda molecule equation with a special boundary condition.

The discrete Toda molecule equation [7] is expressed with the bilinear form as follows

$$\begin{aligned} & \tau_s(m+1, n+1)\tau_s(m, n) - \tau_s(m+1, n)\tau_s(m, n+1) \\ & = q\tau_{s+1}(m, n)\tau_{s-1}(m+1, n+1), \end{aligned} \quad (24)$$

for $s = 1, 2, 3, \dots, N$, q being a constant, with the boundary conditions

$$\tau_0(m, n) = 1, \quad (25)$$

$$\tau_{N+1}(m, n) = f(m)g(n). \quad (26)$$

We replace the boundary condition (26) by

$$\tau_{N+1}(m, n) = F(\tau), \quad (27)$$

in order to obtain the discrete Tzitzeica equation, where $F(\tau)$ is a function of τ_j for $j = 1, 2, \dots, N$.

We have eq.(24) for $N = 2$

$$\begin{aligned} & \tau_1(m+1, n+1)\tau_1(m, n) - \tau_1(m+1, n)\tau_1(m, n+1) = q\tau_2(m, n) \quad (28) \\ & \tau_2(m+1, n+1)\tau_2(m, n) - \tau_2(m+1, n)\tau_2(m, n+1) \\ & = q\tau_3(m, n)\tau_1(m+1, n+1). \end{aligned} \quad (29)$$

Eliminating $\tau_2(m, n)$ of these equations we obtain $\tau_3(m, n)$ expressed by $\tau_1(m, n)$

$$\tau_3(m, n) = q^{-3} \begin{vmatrix} \tau_1(m, n) & \tau_1(m, n+1) & \tau_1(m, n+2) \\ \tau_1(m+1, n) & \tau_1(m+1, n+1) & \tau_1(m+1, n+2) \\ \tau_1(m+2, n) & \tau_1(m+2, n+1) & \tau_1(m+2, n+2) \end{vmatrix}. \quad (30)$$

Hence eq.(30) with the boundary condition

$$\tau_3(m, n) = ab\tau(m+1, n+1)^3/q^3 \quad (31)$$

is the discrete Tzitzeica equation (4) obtained by Schief.

3 A Higher order discrete Tzitzeica equation

The discrete Toda molecule equation (24) suggests that we may extend the discrete Tzitzeica equation (4) to a higher order one. We have for $N = 3$ and for $\tau_0 = 1$

$$\begin{aligned} & \tau_1(m+1, n+1)\tau_1(m, n) - \tau_1(m+1, n)\tau_1(m, n+1) \\ & = q\tau_2(m, n), \\ & \tau_2(m+1, n+1)\tau_2(m, n) - \tau_2(m+1, n)\tau_2(m, n+1) \\ & = q\tau_3(m, n)\tau_1(m+1, n+1), \\ & \tau_3(m+1, n+1)\tau_3(m, n) - \tau_3(m+1, n)\tau_3(m, n+1) \\ & = q\tau_4(m, n)\tau_2(m+1, n+1). \end{aligned}$$

Eliminating τ_2 and τ_3 of these equations we obtain

$$\tau_4(m, n) = q^{-6} \begin{vmatrix} \tau_1(m, n) & \tau_1(m, n+1) & \dots & \tau_1(m, n+3) \\ \tau_1(m+1, n) & \tau_1(m+1, n+1) & \dots & \tau_1(m+1, n+3) \\ \dots & \dots & \dots & \dots \\ \tau_1(m+3, n) & \tau_1(m+3, n+1) & \dots & \tau_1(m+3, n+3) \end{vmatrix}. \quad (32)$$

The *r.h.s.* is a homogeneous polynomial function of τ_1 of degree 4 and is invariant under the following gauge transformation:

$$\tau_1(m, n) \rightarrow \tau_1(m, n) \exp(a_0 m + b_0 n), \quad a_0, b_0 \text{ being constant.}$$

Taking the gauge invariance into account, we propose the following equation

$$\begin{aligned} & \begin{vmatrix} \tau(m, n) & \tau(m, n+1) & \tau(m, n+2) & \tau(m, n+3) \\ \tau(m+1, n) & \tau(m+1, n+1) & \tau(m+1, n+2) & \tau(m+1, n+3) \\ \tau(m+2, n) & \tau(m+2, n+1) & \tau(m+2, n+2) & \tau(m+2, n+3) \\ \tau(m+3, n) & \tau(m+3, n+1) & \tau(m+3, n+2) & \tau(m+3, n+3) \end{vmatrix} \\ &= ab \begin{vmatrix} \tau(m+1, n+1) & \tau(m+1, n+2) \\ \tau(m+2, n+1) & \tau(m+2, n+2) \end{vmatrix}^2, \end{aligned}$$

as an integrable higher order Tzitzeica equation. The integrability of the equation has not been proved yet. However we have a test of identifying integrable discrete systems proposed by Hietarinta and Viellet[8, 9] which analyze the complexity (algebraic entropy) of the map using the growth of the degree of its solution. Numerical calculations of the algebraic entropy of the equation indicate the system's integrability.

4 Ultradiscretization of the discrete Tzitzeica equation

In 1990 one of authors (D.T) and Satsuma reported on a simple cellular automaton which shows soliton-like behavior[10]. It is called "soliton cellular automaton". Tokihiro and coworkers have found that the cellular automaton models are obtained as a special limit, known as the ultradiscrete limit, of the integrable equations[11]. Typically, the limiting procedure simply replaces the summation by the Max operator,

$$\lim_{\varepsilon \rightarrow +0} \varepsilon \log(\exp(A/\varepsilon) + \exp(B/\varepsilon)) = \max(A, B),$$

and the product by the summation,

$$\lim_{\varepsilon \rightarrow +0} \varepsilon \log(\exp(A/\varepsilon) \times \exp(B/\varepsilon)) = A + B.$$

Let

$$V_s(m, n) = \frac{\tau_{s+1}(m, n)\tau_{s-1}(m+1, n+1)}{\tau_s(m+1, n)\tau_s(m, n+1)}. \quad (33)$$

Then the bilinear form

$$\begin{aligned} & \tau_s(m+1, n+1)\tau_s(m, n) - \tau_s(m+1, n)\tau_s(m, n+1) \\ &= q\tau_{s+1}(m, n)\tau_{s-1}(m+1, n+1), \quad \text{for } s = 1, 2, \dots, N, \end{aligned}$$

is rewritten as

$$1 + qV_s(m, n) = \frac{\tau_s(m+1, n+1)\tau_s(m, n)}{\tau_s(m+1, n)\tau_s(m, n+1)}, \quad \text{for } s = 1, 2, \dots, N. \quad (34)$$

Equations (33) and (34) give

$$\frac{V_s(m+1, n+1)V_s(m, n)}{V_s(m+1, n)V_s(m, n+1)} = \frac{[1 + qV_{s+1}(m, n)][1 + qV_{s-1}(m+1, n+1)]}{[1 + qV_s(m+1, n)][1 + qV_s(m, n+1)]}, \quad \text{for } s = 1, 2, \dots, N. \quad (35)$$

The boundary condition $\tau_3(m, n) = ab\tau_1^3(m+1, n+1)$ is transformed, using the expression

$$1 + qV_s(m, n) = \frac{\tau_s(m+1, n+1)\tau_s(m, n)}{\tau_s(m+1, n)\tau_s(m, n+1)},$$

into

$$1 + qV_3(m, n) = \frac{\tau_3(m+1, n+1)\tau_3(m, n)}{\tau_3(m+1, n)\tau_3(m, n+1)} = [1 + qV_1(m+1, n+1)]^3.$$

Accordingly we have the discrete Tzitzeica equation in the ordinary form,

$$\begin{aligned} \frac{V_1(m+1, n+1)V_1(m, n)}{V_1(m+1, n)V_1(m, n+1)} &= \frac{[1 + qV_2(m, n)]}{[1 + qV_1(m+1, n)][1 + qV_1(m, n+1)]}, \\ \frac{V_2(m+1, n+1)V_2(m, n)}{V_2(m+1, n)V_2(m, n+1)} &= \frac{[1 + qV_1(m+1, n+1)]^4}{[1 + qV_2(m+1, n)][1 + qV_2(m, n+1)]}. \end{aligned}$$

Following the limiting procedure we put

$$V_s(j, k) = \exp(x_s(j, k)/\epsilon), \quad q = \exp(-c/\epsilon), \quad \text{for } s = 1, 2, 3.$$

The discrete Tzitzeica is transformed into the following max-plus equations in the small limit of ϵ ,

$$\begin{aligned} x_1(m+1, n+1) &= x_1(m+1, n) + x_1(m, n+1) - x_1(m, n) \\ &\quad + \max(0, x_2(m, n) - c) - \max(0, x_1(m+1, n) - c) \\ &\quad - \max(0, x_1(m, n+1) - c), \\ x_2(m+1, n+1) &= x_2(m+1, n) + x_2(m, n+1) - x_2(m, n) \\ &\quad + 4 \max(0, x_1(m+1, n+1) - c) - \max(0, x_2(m+1, n) - c) \\ &\quad - \max(0, x_2(m, n+1) - c). \end{aligned}$$

We remark that the discrete sine-Gordon equation is obtained in connection with a discrete geometry [12] as well as the discrete Tzitzeica equation. However it is very difficult to ultradiscretize the former equation in contrast to the latter.

5 Periodic Boundary Conditions

We transform the coordinates m, n into new coordinates j, k :

$$m = \frac{1}{2}(j+k), \quad n = \frac{1}{2}(j-k),$$

in order to obtain equations with the periodic boundary condition. We express functions of m, n in the new coordinates

$$\begin{aligned} V_s(m, n) &= V_s(j, k), & V_s(m+1, n) &= V_s(j+1, k+1), \\ V_s(m, n+1) &= V_s(j+1, k-1), & V_s(m+1, n+1) &= V_s(j+2, k). \end{aligned}$$

Then, the discrete Tzitzeica equation is transformed into

$$\begin{aligned} &\frac{V_1(j+2, k)V_1(j, k)}{V_1(j+1, k+1)V_1(j+1, k-1)} \\ &= \frac{[1+qV_2(j, k)]}{[1+qV_1(j+1, k+1)][1+qV_1(j+1, k-1)]}, \end{aligned} \quad (36)$$

$$\begin{aligned} &\frac{V_2(j+2, k)V_2(j, k)}{V_2(j+1, k+1)V_2(j+1, k-1)} \\ &= \frac{[1+qV_1(j+2, k)]^4}{[1+qV_2(j+1, k+1)][1+qV_2(j+1, k-1)]}. \end{aligned} \quad (37)$$

It is generally accepted that the system's integrability is not destroyed by imposing a periodic boundary condition on the system. We use the periodic boundary condition $V_s(j, k) = V_s(j, k+N)$ in order to transform the partial difference equations (36) and (37) into a coupled form of ordinary difference equations. The algebraic entropy of the higher order discrete Tzitzeica equation (33) is calculated by imposing the periodic boundary condition $V_s(j, k) = V_s(j, k+N)$ with periods $N = 1, 3, 5, \dots$.

The discrete Tzitzeica equation is reduced, for $N = 1$ (uniform in k -direction), to

$$\frac{V_1(j+2)V_1(j)}{V_1^2(j+1)} = \frac{[1+qV_2(j)]}{[1+qV_1(j+1)]^2}, \quad (38)$$

$$\frac{V_2(j+2)V_2(j)}{V_2^2(j+1)} = \frac{[1+qV_1(j+2)]^4}{[1+qV_2(j+1)]^2}. \quad (39)$$

Equations (38) and (39) give a relation

$$\left[\frac{V_1(j+3)V_1(j+1)}{V_1^2(j+2)} \right]^2 \frac{V_2(j+2)V_2(j)}{V_2^2(j+1)} = 1.$$

Hence

$$V_2(j) = c_0 c_1^j V_1^{-2}(j+1), \quad (40)$$

where c_0 and c_1 are integration constants. Substituting the relation into eq.(38) we obtain

$$\frac{V_1(j+2)V_1(j)}{V_1^2(j+1)} = \frac{[1 + qc_0c_1^j V_1^{-2}(j+1)]}{[1 + qV_1(j+1)]^2}. \quad (41)$$

Equation (41) is a non-autonomous equation if $c_1 \neq 1$, while the Tzitzeica equation is an autonomous equation. Hence we choose $c_1 = 1$. Putting $V_1(j+1) = x_n$ and $c_0 = q = c$ we obtain a one-dimensional discrete Tzitzeica equation in the following form,

$$x_{n+1}x_{n-1} = c^2 \frac{(1 + x_n^2/c^2)}{(1 + cx_n)^2}. \quad (42)$$

Let $x_n = e^{X_n/\epsilon}$, $c = e^{C/\epsilon}$. Then the one-dimensional discrete Tzitzeica equation (42) is transformed into the following max-plus equations in the small limit of ϵ ,

$$X_{n+1} = 2C + 2 \max(0, X_n - C) - 2 \max(0, X_n + C) - X_{n-1}. \quad (43)$$

6 Periods of the ultradiscrete Tzitzeica equation

Equation (43) has the following conserved quantity,

$$H = \max(X_{n-1} - X_n - 2C, -X_{n-1} + X_n - 2C, -X_{n-1} - X_n, X_{n-1} + X_n), \quad (44)$$

where H is constant for n . This quantity H means that eq.(43) is integrable. Moreover, a solution from any initial data has the following remarkable features;

- (a) If $|(H - 2C)/4C|$ is rational, the solution is periodic with a finite period. Otherwise, the period becomes infinite.
- (b) The finite period of a solution is determined only by C and H .
- (c) Therefore, if C is fixed, finite periods of all solutions with the same H are the same.

If we consider eq.(43) is a mapping from (X_{n-1}, X_n) to (X_n, X_{n+1}) , we can show the above features by a geometrical analysis of solution orbits in a phase plane. We omit the details of the analysis since it needs a long space and only show how a period is determined by C and H . Consider the following relation,

$$\frac{q}{p} = \left| \frac{H - 2C}{4C} \right|, \quad (45)$$

where p and q are positive integer and relatively prime. Then the period is expressed by p and q as follows.

$C > 0$	$H = 0$	2
	$0 < H < 2C$	$3p - 4q$
	$H = 2C$	3
	$2C < H$	$3p + 4q$
$C = 0$	$0 \leq H$	4
$C < 0$	$H = 2 C $	1
	$2 C < H$	$-3p + 4q$

Note that $0 \leq H$ if $0 \leq C$ and $2|C| \leq H$ if $C < 0$, which can be shown from eq.(44).

For example, if $C = 3$, $X_0 = 1$ and $X_1 = 3$, $H = 4$ and $|(H - 2C)/4C| = 1/6$. Then we have $q = 1$ and $p = 6$, and the period becomes $3p - 4q = 14$. Indeed, the solution becomes

$$X_n : 1, 3, -7, 3, 1, -5, 5, -1, -3, 7, -3, -1, 5, -5, 1, 3, \dots$$

If we take the same C and $X_0 = X_1 = 2$, the same H , p and q are obtained. The solution is

$$X_n : 2, 2, -6, 4, 0, -4, 6, -2, -2, 6, -4, 0, 4, -6, 2, 2, \dots,$$

and the period is also 14.

7 Concluding remarks

We have ultradiscretized the discrete Tzitzeica equation by Schief. The discrete Tzitzeica equation is connected to the discrete affine geometry. So it would be of strong interest to find a geometry connected to the ultradiscrete Tzitzeica equation.

References

- [1] Tzitzeica G., *C.R. Acad.Sci*, T.150, (1910) pp.955-956.
- [2] Tzitzeica G., *C.R. Acad.Sci*, T.150, (1910) pp.1277-1229.
- [3] O.V.Kaptsov and Yu.V.Shan'ko, Trilinear representation and the Moutard transformation for the Tzitzeica equation, *solv-int/9704014* v1 21 Apr 1997.
- [4] W.K.Schief, Self-dual Einstein spaces via a permutability theorem for the Tzitzeica equation, *Phys.Lett. A* 223 (1996) pp.55-62.
- [5] W.K.Schief, Self-dual Einstein spaces and a discrete Tzitzeica equation, a permutability theorem link, in *Symmetries and Integrability of Difference Equations*, Edited by P.A. Clarkson and F.W. Nijhoff, London Mathematical Society Lecture Note Series 225 (1999) pp.137-148.
- [6] Ryogo Hirota, *The Direct Method in Soliton Theory* (in Japanese), (Iwanami, 1992).
- [7] Ryogo Hirota, Discrete Two-Dimensional Toda Molecule Equation, *J. Phys. Soc. Jpn.* **56**(1987) pp.4285-4288.
- [8] J. Hietarinta and C-M. Viallet, Singularity Confinement and Chaos in Discrete Systems, *Phys.Rev. Lett.* **81**(1998) pp.325-328.
- [9] M.P. Bellon and C-M. Viallet, Algebraic entropy, *chao-dyn/9805006* v3 3 Jun 1998.
- [10] D.Takahashi and J.Satsuma, A Soliton Cellular Automaton, *J. Phys. Soc. Jpn.* **59**(1990) pp.3514-3519.
- [11] T.Tokihiro, D.Takahashi, J.Matsukidaira and J.Satsuma, From Soliton Equations to Integrable Cellular Automata through a Limiting Procedure, *Phys.Rev.Lett.* **76**(1996) pp.3247-3250.
- [12] A.I. Bobenko and U. Pinkall, Discrete surfaces with constant negative Gaussian curvature and the Hirota equation *J.Diff.Geom.* **43** (1996) pp.527-611.